

Serum Cholesterol & Chronic Low Back Pain!


Written by: Dr. Bahram Jam, PT

Advanced Physical Therapy Education Institute (APTEI), Thornhill, ON, Canada
August 5, 2009 Article published on www.aptei.ca "Clinical Articles"

There have thousands of papers, research studies on the mysterious condition referred to as non-specific chronic low back pain (LBP). As physiotherapists, we are all well aware of the numerous potential sources and contributing factors to LBP, which include various patho-anatomical and psychosocial causes. However, one contributing factor that many health care providers including physiotherapist may often fail to recognize, is the influence of serum cholesterol on LBP. The objective of this short article is to summarize a few of the research papers on the topic of atherosclerosis, cholesterol and LBP.

Several papers have been written on the association between trans fats and a wide range of diseases including Alzheimer's disease, coronary heart disease, prostate cancer and obesity¹. There is also little debate that trans fats increase LDL cholesterol (the bad cholesterol) and decrease HDL (the good cholesterol)¹. Now the question is, is there an association between trans fats, LDL cholesterol and LBP?

The branching arteries of the abdominal aorta, including the four paired lumbar arteries and the middle sacral artery feed the lumbar spine (Fig. 1). Atherosclerosis in the wall of the abdominal aorta may block the relatively small orifices of lumbar and middle sacral arteries. Obstruction of these arteries inevitably leads to ischemia in the lumbar spine^{2,3}.



It has been suggested that the reduced blood flow into the intervertebral discs, vertebral bodies and myofascial structures could result in various back symptoms².

The aim of this recent systematic literature review² was to evaluate the links between atherosclerosis and degenerative disc disease (DDD) or LBP. Following a Medline/PubMed database search for all published articles on atherosclerosis and DDD/LBP, 179 papers were identified. The search was performed with the medical subject headings atherosclerosis, cardiovascular risk factor, or vascular disease and keywords "disc degeneration", "disc herniation", and "back pain". After the exclusion of low quality studies, 25 papers were included.

The 6 basic findings of this systematic review were:

1: Post-mortem studies showed an association between **aortic atherosclerosis and DDD**.


2: Post-mortem studies showed a strong association between **occluded lumbar arteries and a life-time of LBP**.

3: Clinical studies showed that **aortic calcification was associated with LBP**.

4: Clinical studies showed that **stenosis of lumbar arteries was associated with both DDD and LBP**.

5: Epidemiological studies showed that **smoking and high serum cholesterol levels were the most consistent associations with DDD and LBP**.

6. Cohort large studies showed clear **associations between elderly people with cardiovascular risk factors and LBP**.



Here is a summary of one sample paper published in the journal 'Spine' which was included in the above-mentioned systematic review 3. Magnetic resonance (MR) aortography and LDL cholesterol blood tests were performed on patients with persistent non-specific LBP. The patients ranged from 35 to 70 years of age (mean of 56 years). The 4 basic findings of this study were:

1: Over 75% (that's 3 out of 4 patients) of both the men and women showed **occluded lumbar and/or middle sacral arteries**.

2: The prevalence of occluded lumbar arteries was **2.5 times more** in the LBP patients than the age matched control group.

3: Disc degeneration was significantly associated with occluded lumbar/middle sacral arteries.

4: Patients with **higher serum LDL cholesterol** levels had significantly greater neurogenic symptoms and complained more often of severe pain than those with normal LDL cholesterol.

Clinical Relevance / Personal Comment:

Firstly, is it not impressive that there are this many studies published in peer reviewed medical journals on this rarely spoken topic... cholesterol, atherosclerosis, DDD and back pain? With so much focus on patho-anatomical and psychosocial causes of LBP, vascular disease as a contributing factor to chronic, non-mechanically responsive LBP has been regrettably undermined. Inevitably patients with lumbar and sacral artery atherosclerosis fail to respond to NSAIDs, extension exercises, manual therapy, modalities, traction, acupuncture, stabilization, etc. It is also probable that some patients with 'failed back surgery' may have had a coincidental disc herniation, but an underlying deficient lumbar vascular supply as the primary contributing factor to their LBP and 'sciatica'. The ever so widely accepted idea that DDD is inevitable and is simply due to 'old age' may in fact not be fully accurate. Perhaps by controlling atherosclerosis through proper medical care, stress management, nutrition and exercise, the progression of DDD can be controlled.

Longitudinal studies are still not available to support this hypothesis.

Secondly, considering the direct association between cardiovascular disease, high LDL cholesterol and LBP, every patient with non-mechanically responsive persistent LBP should be questioned about their cardiovascular health. Therefore, clinicians should ideally ask the following questions to see if diet, cholesterol and atherosclerosis are potential contributing factors to a patient's LBP.

- Do you know if you have significant stress?
- Do you have hypertension / high blood pressure?
- Do you smoke?
- Do you have a history of heart disease?
- Do you do any aerobic or physical exercises at least 3 times per week?
- Do you eat at least 2 servings of fresh fruits everyday?
- Do you eat at least 3 servings of fresh vegetables everyday?

Clinical Management Options:

- Manage stress, as it has been hypothesized to be a primary cause of atherosclerosis. Consider seeking professional help or seriously changing occupation or lifestyle.
- Consult a Registered Dietician or a naturopathic doctor for a nutritional evaluation

- Avoid or at least reduce the consumption of processed grains, sugars, high fructose corn syrup
- Avoid or at least reduce processed animal fat consumption
- Consider a Mediterranean type of diet, consisting of daily fresh fruits and vegetables, virgin olive oil and fish
- Consider eating fish more often and/or take fish oil capsules (for Omega 3s)
- With the guidance of a physiotherapist, start a gentle yet progressive aerobic exercise program ...anything for 10-30 minutes at least 5X/week
- Stop smoking

My intention for sharing this information with fellow physiotherapists is not to undermine the importance of physiotherapy intervention for a sub-group of patients with chronic LBP and DDD; in fact it is the opposite. Based on hundreds of clinical trials on the topic of exercise and serum cholesterol levels, the value of a regular aerobic exercise program for individuals with cardiovascular disease cannot be over emphasized.



Regrettably, the primary medical intervention and focus for the management of high LDL cholesterol and hypertension continues to be only a pharmaceutical approach⁴. Three separate meta-analysis studies examining the effects of aerobic exercise on lipids and lipoproteins have concluded that regular aerobic exercise is efficacious for increasing HDL cholesterol and decreasing LDL cholesterol, and triglycerides⁵⁻⁷. Exercise has been shown to

be even more effective in subjects with initially high total cholesterol levels or low body mass index⁵.

As physiotherapists, we are the “exercise specialists” with the most favourable educational training to provide an effective and patient specific exercise prescription for individuals presenting with either LBP or cardiovascular disease and in some cases both.

References:

1. Stender S, et al Fast food: unfriendly and unhealthy. *Int J Obes (Lond)*. 2007 Jun;31(6):887-90. Epub 2007 Apr 24
2. Kauppila LI. Atherosclerosis and Disc Degeneration/Low-Back Pain -A Systematic Review. *Eur J Vasc Endovasc Surg*. 2009 Mar 25.
3. Kauppila LI, Mikkonen R, Mankinen P, Pelto-Vasenius K, Mäenpää I. MR aortography and serum cholesterol levels in patients with long-term nonspecific lower back pain. *Spine*. 2004 Oct 1;29(19):2147-52.
4. Evans M, Roberts A, Davies S, Rees A. Medical lipid-regulating therapy: current evidence, ongoing trials and future developments. *Drugs*. 2004;64(11):1181-96.
5. Kodama S, Tanaka S, Saito K, Shu M, Sone Y, Onitake F, Suzuki E, Shimano H, Yamamoto S, Kondo K, Ohashi Y, Yamada N, Sone H. Effect of aerobic exercise training on serum levels of high-density lipoprotein cholesterol: a meta-analysis. *Arch Intern Med*. 2007 May 28;167(10):999-1008.
6. Kelley GA, Kelley KS, Tran ZV. Aerobic exercise and lipids and lipoproteins in women: a meta-analysis of randomized controlled trials. *J Womens Health (Larchmt)*. 2004 Dec;13(10):1148-64
7. Kelley GA, Kelley KS, Franklin B. Aerobic exercise and lipids and lipoproteins in patients with cardiovascular disease: a meta-analysis of randomized controlled trials. *J Cardiopulm Rehabil*. 2006 May-Jun;26(3):131-9; quiz 140-1 Pattanittum P1, Turner T, Green S, Buchbinder R. Non-steroidal anti-inflammatory drugs (NSAIDs) for treating lateral elbow pain in adults. *Cochrane Database Syst Rev*. 2013 May 31;5:

Study summaries are available on www.aptei.ca “Clinical Articles” accessible by APTEI
Report subscribers
To request a copy of this article simply email jam@aptei.com